

**Republic of Iraq
Ministry of Higher Education
& Scientific Research
University of Al-Qadisiyah
College of Veterinary Medicine**



Chemical contamination of meat

A Graduation Project Submitted to the College of Veterinary Medicine/ University of Al-Qadisiyah in a partial fulfillment of the requirements for the Degree of Bachelor of Science in Veterinary Medicine and Surgery.

By
Sarah Mohsen Rahim

Supervised by
Afrah Sabeeh Mhyson

2021 A.D.

1441 A.H.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

فَنَعَلَى اللَّهِ الْمَلِكُ الْحَقُّ وَلَا تَعْجَلْ بِالْقُرْآنِ مِنْ قَبْلِ أَنْ يُقْضَىٰ
إِلَيْكَ وَحْيُهُ، وَقُلْ رَبِّ زِدْنِي عِلْمًا ﴿١١٤﴾

صَدَقَ اللَّهُ الْعَظِيمُ،

من سورة طه

Certificate of Supervisor

I certify that the project entitled was prepared by **Sarah Mohsen Rahim**
under my supervision at the College of Veterinary Medicine / University of
Al-Qadissiya.

Assis.Prof.Afrah Sabeeh Mhyson

Dept. of Public Health

Coll. Of Vet. Med./ Univ. of Al-Qadisiyah

Certificate of Department

We certify that as finished her Graduation Project entitled

Instructor

Dr. Muthanna H. Hussain

Head of Dept. of Int. and Prev. Med.

Dr. Muthanna H. Hussain

Acknowledgment

Great truth of God My God is not the night except with your thankfulness and the daytime to your obedience. To the Prophet of mercy and the light of the worlds .. Prophet Muhammad peace be upon him I am extremely grateful to the Ministry of Higher Education and University of Al-Qadisiyah Collage of Veterinary Medicine Department of Public Health , and deep respect to my supervisor Dr. **Afrah Sabeeh Mhyson**. for her patience, motivation, immense knowledge and continuous support in the course of this project.

List of content

No.	Subject
1.	Introduction
2.	Literatures Review
2.1.	Pesticides
2.2.	Heavy metals
2.3.	Veterinary drugs
2.4.	Antibiotics
2.5.	Anabolic compounds
2.6.	Melamine
2.7.	Mycotoxins
3.	Conclusion
4	Recommendations
5	References

Abstract

Humans around the world are exposed to chemical contaminants during their life time. among the thousands of existing contaminates, some are persistent and remain in the environment for years. The variation in measurable levels depends mainly on the fact that some are synthesized as industrial products, whereas others are released accidentally, as by-products, or given to animals as growth promoters or as prophylactic or therapeutic agents. The measurement of these contaminants requires a complex procedure including sample extraction, sample clean-up, and physico-chemical analysis after chromatographic separation. Contaminants such as organochlorine pesticides, heavy metals, microbes, melamine, hormones, antibiotic, and other feed additives are often measured in various types of matrices during food safety programs, environmental monitoring, and epidemiological studies. Serious health problems including cancer, kidney diseases, disarray and other diseases in humans might be related to food contaminates. According to the World Health Organization (WHO), 1.8 million people died from diarrhea related diseases in 2005. Children and developing fetuses are generally at greater risk from exposure to different chemicals. A great number of these cases might be attributed to contaminate found on food. More than 90% of human exposure to harmful materials is due to consumption of contaminated food items such as meat, milk and dairy products, as well as fish and derived products. The overwhelming majority of food poisoning in the world might originate from contaminated animal products and there are millions of cases, and hundreds of deaths, caused by contaminated animal products every year. This paper will discuss and highlight the most common chemical and biological contaminates in meat and meat products from various species which threaten human health. Key words: Meat; meat products; contaminants; human health.

1.Introduction

The principle contaminants of food-stuff in the world include microorganisms and their metabolites, mycotoxins, heavy metals, nitrates, nitrites, hormones, pesticide residues, antibiotics, dioxins, polychlorinated biphenyls, genetically modified organisms, Toxic pigments, and melamine. Meat and meat products are integral parts of the daily diet for many people due to tradition, variety, reasonable prices, versatility, satiety value and Taste. It is the nutritional value of meat, however, which distinguishes it as a beneficial component in a balanced diet [1]. Meat provides a ready source of protein, minerals and Some vitamins (B6 and B12). Meat is a rich source of many nutrients per unit energy; Therefore, emphasis should be placed on the importance of meat safety for human consumption. traditional meat inspection systems are to a large extent based on the Concept of inspecting live animals (ante-mortem) and carcasses (post-mortem) to certain the presence of items hazardous to human health [2]. Judgment of gross findings relative to the suitability of the product for human consumption is made according to Standardized criteria. However, traditional methods of meat and fish inspection do not determine contaminants, such as antibiotics, anabolic agents, melamine, and heavy metals [3]. The objective was to review the current situation of meat and meat products contaminants with residues of hormones, antibiotics, pesticides, melamine, and heavy metals in meat and fish products. moreover, pilot studies were carried out to detect hormones, antibiotics, heavy metals and melamine in meat and fish collected from local market. samples were collected from the major supermarket outlets and butcher shops and evaluated ELISA.

2.Literatures Review

2.1. Pesticides

more than 100 pesticides are known, but it is unlikely to find all of them in one food-stuff. They are mainly used in agriculture (68%), commercial and industrial activities (17%), domestic settings (8%) and in governmental applications (7%) [4] The widespread presence of organochlorine pesticide residues (OCPs) in Food has arisen due to their extensive agricultural application and industrial emissions in The environment. A large number of pesticides may potentially be used in the production of agricultural crop commodities, leading to indirect exposure of animals through feed and the potential for residues in animal products [5] Exposure to pesticides can occur through ingestion of crops or crop by-products that are Used as animal feed. significant animal exposure to pesticides usually occurs through the Intake of grain, fodder and forage, as well as feeding on processed by-products, meals and waste. Animals may also be exposed to pesticides by ingestion of contaminated soil While grazing. Less frequent sources of livestock exposure to pesticides include Accidental spills, improper waste disposal and contaminated areas from past use of products that are persistent in the environment [6] many pesticides tend to accumulate in animal tissues and are a function of the physicochemical properties of the pesticides [7]. The lipophilic compounds have a great tendency to accumulate in animal tissues and products [8]. One reason for the environmental persistence of these compounds is that microorganisms are either Unable to degrade them or do so at a very slow rate [9]. persistent organic Chlorine compounds such as dichlorodiphenyltrichloroethane (DDT) play an important Role in chronic poisoning and contribute to a number of pathological processes [10] Fat solubility of these compounds is Responsible for their varied concentrations in tissues and their accumulation in the Lipoproteins of the cell membranes, thus changing their structure and permeability[11]However, compounds that are Water-soluble or can be readily metabolized to water-soluble products are rapidly Excreted in the urine and do not give rise to significant residues in meat. pesticides can be taken up by livestock in their feed or water and may be Incompletely eliminated at the time of slaughter [12].

These compounds tend to accumulate in body tissues. They are resistant to chemical or biological degradation and persist in the environment for a considerable period of time. They can be absorbed by all routes, including inhalation, ingestion and Dermal absorption. The awareness that OPPs may also concentrate along the food chain Has led to the establishment of low maximum residue limits (MRLs) in meat and fish. consequently, this requires the control of this type of compound in fatty matrices [13]. pesticides may induce or aggravate certain health problems in Humans such as cancer, immune system disturbances and disruption of hormonal Functions [14] Residues of OCPs were found in meat and meat products [15]at different levels depending on the type of pesticide and meat. Although DDT had been banned, however, it is a widely used insecticide that persists in animal tissues. other cyclodiene groups also tend to accumulate in body fat [16]. The accumulation of pesticides in the liver may be as much as 100 times higher [16] lactating cows excrete a considerable amount of these compounds in the milk, and residues are also found in eggs of laying hens. The intermuscular and Intramuscular fat determines to a great extent the amount of these compounds ingested by the consumer. Polychlorinated biophenyls (PCBs) are synthetic compounds with high Chemical and thermal stability. PCBs are extremely persistent and soluble in body tissues and have tumor-promoting effects [16]

2.2. Heavy metals

Anthropogenic activities may have an adverse impact on human health due to The discharge of industrial waste and domestic sewage [17].

These effluents usually contain elevated levels of heavy metals that accumulate in macro-Particles which form the basis of many food chains [18] non-essential elements such as lead, cadmium and mercury, have toxic properties and a tendency to accumulate in meat and fish products. Significant Concentrations of these elements were found in edible animal tissue such as muscle, liver and kidney. Eight per cent of the lead in food is taken up from the human gastrointestinal Tract. Lead interferes with haemoglobin synthesis and is able to inhibit several enzymes [19].

Cadmium accumulates in the kidneys and it has an extremely long Biological half-life in humans in the order of 20-30 years [20]. five to ten per Cent of cadmium is absorbed and bound in the liver with metallothionein and stored in the Kidney as Cd-thionein. kidneys which contain more than 200 mg cadmium per kg may exhibit morphological

abnormalities in the tissue structure and increase excretion of Proteins, amino acids, glucose and calcium [21]. other symptoms of cadmium poisoning are emphysema of the lungs, anemia and insufficient bone mineralization as a Result of increased calcium excretion. Cadmium concentrations in old animal tissues are Sometimes extremely high. Mercury levels above 0.05 mg/kg in meat and fish products Should be considered harmful or hazardous. Copper and thallium are also elements that warrant concern [22].

2.2.Veterinary drugs

Veterinary drugs are generally used in farm animals for therapeutic and Prophylactic purposes. They include a large number of different types of compounds Which can be administrated in the feed or in the drinking water. The residues may also originate from contaminated animal feedstuffs [23]. The presence of residues and its associated harmful health effects on humans makes the control of veterinary drug Residues an important measure in ensuring consumer protection. Residues include Anabolic compounds and hormones in animal products.

2.3.Antibiotics

Antibiotics contaminating livestock products (milk, eggs and meat) are dangerous because they can induce allergic reactions, disturb the gastrointestinal microflora. previous studies at Sultan Qaboos University demonstrated that various types of antibiotic residues were found in meat samples of poultry, sheep and goats [24]. anti-microbial activity has also been found in all commercial milk products sold in Oman [25]. antibacterial drugs are usually used in animals for therapeutic, preventative and nutritive purposes [26]. If proper withdrawal times are not Adhered to, residues may be present in the animal tissues at slaughter. the use of antibiotics for growth promotion or prevention and treatment of infectious diseases, can result in resistance among both resident bacterial pathogens and commensal organism [27] which can cause allergic reactions and transfer of resistance to humans [28].antibiotic resistance is most clearly a problem when it leads to therapeutic failure or the need to use more costly, toxic, or expensive drugs [29] and it can also be a problem when it increases the frequency, duration or severity of infection[30].

In humans, microbial resistance is an important problem in a wide variety of Infections of the skin, respiratory, genito-urinary and gastrointestinal tract [31]. there is considerable evidence that antibiotic used in animals, caused the selection of resistant strains that can then spread to humans [32].

2.4. Anabolic compounds

Anabolic compounds anabolic agents influence the metabolic functions of animals resulting in an increased growth rate and also an increased deposition of proteins in the muscle mass. A primary effect is an increase in the protein deposition, usually linked to fat utilization that decreases the fat content in the carcass and increases meat leanness [33]. many of the anabolic compounds have oestrogenic or androgenic properties. The residues of anabolic agents or their metabolites in fish, meat and other foods of animal origin may cause adverse toxic effects on consumers' health. It has been shown that there was an association between some forms of hormone-dependent cancers and red meat consumption [34]. the consumption of lamb and beef meat containing residues of clenbuterol was associated with symptoms in humans characterized by gross tremors of the extremities, tachycardia, nausea, headaches and dizziness.the implantation of 17β -estradiol in meat animals results in high concentrations in edible tissue. Estradiol is carcinogenic in reproductive, bone, pituitary and lymphoid tumors. Thyreostats, which have been prohibited in many countries, are inhibiting the thyroid function which allows a significant increase in the live weight of animals due to water retention in muscle mass. Residuals of this hormone may be found in muscle tissues and can interfere with normal hormonal action [35].

It has been reported that high doses of progesterone administered over a long period of time are correlated with an increase in the rate of tumor growth [36] .It stimulates cell division in hormonally sensitive tissues, thereby increasing the possibility for accumulation of random errors during DNA duplication [37] .This increased cell proliferation also has the effect of stimulating growth of mutant cells [38] .Some women taking estrogen supplements develop breast or uterine cancer, indicating that estradiol may be one of several factors important in the development of these cancers [39]Two outbreaks, one of breast enlargement in young school children [40] and another of precocious sexual development [41]were suggestive of exposure to environmental estrogenic compounds, possibly Zeranol. The researchers suspected that one consignment of meat might have contained residues of some estrogenic compounds

and reported that symptoms gradually disappeared after the children stopped consuming meat. Carcinogenicity studies of trenbolone in rodents indicated that 10 or 100 ppm ($\mu\text{g/g}$) trenbolone in the diet was associated with an increase in liver cancer in mice and 50 ppm trenbolone may have enhanced pancreatic cancer in rats [42]. The β_2 – agonists (adrenaline) are produced by the adrenal gland and are released when an animal is under stress. synthetic compounds (β -adrenergic agonists) have been shown to repartition nutrients from fat to muscle. the doses of β_2 –agonists necessary to cause repartitioning are 5-10 times greater than the recommended therapeutic doses. there is a potential risk to consumer's health at the high doses necessary for growth promotion. There have been documented cases of mass intoxications after consumption of liver and meat treated with Clenbuterol. Use of the highly active β_2 –agonists as growth promoters is not appropriate because of the potential hazard for human and animal health.

2.5.Melamine

Melamine (2,4,6-triamino-1,3,5-triazine) is an organic compound commonly used to produce various products, including dishes, plastic resins, in fertilizer, as a flame retardant, manufacture of wrinkle-free textiles and components of paper and paperboard that may come into contact with meat [43] .the researchers used melamine as a non-protein nitrogen source for feeding animals and concluded that the melamine was digested but most of it was eliminated because micro flora in the rumen was not able to assimilate. As a result, trace amounts of melamine and its three analogues, cyanuric acid, ammelide and ammeline, referred to as melamine compounds may be present in animal tissue. The average concentration of melamine in food from approved industrial uses is estimated to be less than 0.015 parts per million [44]. these levels of melamine in food are extremely minute and do not pose a public health concern. Intentional addition of melamine to food, however, does pose a significant risk. these levels of melamine and related analogues in food from accepted uses that do not result from adulteration or misuse. this includes expected levels from the environment, food processing, packaging materials, residues from the legitimate use of triazine pesticides or veterinary drugs, and legitimate use of melamine in fertilizers or cyanuric acid in feed additives. however, the intentional addition of melamine and/or analogues directly to food, food ingredients, animal feed, feed ingredients or pelletizing agents is not acceptable. It may also be present indirectly in foods of animal origin as a result of

carryover from the intentional addition to animal feed, therefore, there is an urgent need to detect melamine in foods [45]. In particular, it was discovered that melamine-contaminated ingredients had been used to prepare feed for chickens and fish. A preliminary study at Sultan Qaboos University demonstrated that various levels of melamine were found in poultry meat (Unpublished data).

2.7. Mycotoxins

One of the most dangerous pollutants of food-stuff is fungi and their metabolites – mycotoxins, which can cause mycotoxicosis. Mycotoxins are found on many plant products, particularly oilseeds and cereals. These substances are characterized by acute and chronic toxic properties, while some of them are potent carcinogens. Although animals are an effective toxin eliminator [46], residues of mycotoxin can still be present in animal tissue. The carcinogenic potential of aflatoxins B1 is the highest. Aflatoxins are stable, heat resistant compounds and can be found in animal tissue [47].

3. Conclusion

The chemical contamination of food has emerged as a serious concern with potential health hazards in their wake. Majority of the food contamination occurs through naturally occurring toxins and environmental pollutants or during the processing, packaging, preparing, storage, and transportation of food. As the technology advances, the detection of such contaminants becomes easier. However, there are several contaminants that are still unknown and research continues in this regard. Although the government has taken adequate steps to minimize the individual exposure to food contaminants, there are still measures that need to be taken to reduce the health risks and diseases that come with the chemical food contamination.

4. Recommendations

There is legislation in place to regulate the levels of several chemicals in the food. Unhealthy additives and adulterants are legally not allowed for use. However, effective surveillance and response systems are required to prevent chemical hazards from entering the food supply and posing harm to the public. The minimum levels of chemicals that are allowed in food, such as pesticide concentration should not go higher than the limit assigned. However, errors may still occur in following the determined concentration and guidelines. Particularly in the case of developing and

underdeveloped countries, the legislation enforcement is still weak about administrating the concentration of harmful contaminants in the food. Some countries are highly dependent on agriculture, resulting in high levels of pesticides seeping into the ground water, contaminating both food and water. Non-regulated chemicals are of specific concern and more research needs to focus on contaminants that escape human detection. Also, individual consumer concerns are essential as they can play a fundamental role in managing their health. Moreover, the popularity and widespread use of internet also allow consumers to seek information online and reduce the health risks associated with food contamination incidents. The news media and journalists have an important role in reporting on the outbreaks, threat and its cause, including expert commentary regarding the chemical food contaminants. Furthermore, the public need to keep a healthy degree of skepticism about the contaminated food products reported on the news and avoid consuming the accused food products until scientific evidence justifies immediate action. Most importantly, the food industries must accept the need to be more honest and upfront in producing safe commercial food products as well as protecting the public from food contamination.

5.References

1-Abou-Arab, A. A. K. (2002). Degradation of organochlorine pesticides by meat starter in liquid media and fermented sausage. *Food and Chemical Toxicology*, 40: 33-41.

2-Addison, R. F., Brodle, P. F., and Zinck, M. E. (1984). DDT has declined more than PCBs in eastern Canadian seals during the 1970s. *Environmental Science and Technology* 18: 935-937.

3-Akkoyum, A., Kohen, V. F., and Bilitewski, U. (2000). Detection of sulphamethazine with an optical biosensor and anti-idiotypic antibodies. *Sensors and Actuators B*, 70, 12- 18.

4-Alexander, M. (1981). Biodegradation of chemicals of environmental concern. *Science*, 211: 132-138.

5-Alloway, B. J. and Ayres, D. C. (1997). *Chemical Principles of Environmental pollution*. Chapman and Hall, London.

6-Antunes-Madeira, M. C., Almeida, L. M., and Madeira, V. M. C. (1993). Depth-dependent effects of DDT and lindane on the fluidity of native membranes and extracted lipids implication for mechanisms of toxicity. *Bulletin of Environmental contamination and Toxicology*, 5: 787-794.

7-Arino, A., Herrera, A., Conchello, M. P., and Perez, C. (1992). Hexachlorobenzene residues in Spanish meat products after cooking, curing and long-term ripening. *Journal of Food Protection*, 55: 920-923.

8-Arino, A., Herrera, A., Conchello, M.P., Lazaro, R. and Perez, C. (1993). Hexachlorobenzene residues in meat products after processing. *Journal of Food Composition Analysis*, 6: 55-61.

9-Arino, A., Lazaro, R. Conchello, M. P., Bayarri, S., and Herrera, A. (1995). The effect of processing on incurred residues on DDE in meat products. *Food Additives and Contaminants*, 12: 559-566.

10-Barbosa, J. Cruz, C., Martins, J., Silva, J. M., Neves, C. and Alves, C. (2005). Food poisoning by clenbuterol in Portugal. *Food Additives and Contaminants*, 22: 563- 566.

11-Barza, M. (2002). Potential mechanisms of increased disease in human from antimicrobial resistance in food animals. *Clinical Infectious Diseases*, 34:123- 125.

12-Belloque, J., Garcia, M. C., Torre, M. and Marina, M. L. (2002). Analysis of soybean proteins in meat products: A review. *Critical Reviews in Food Science and Nutrition*, 42: 507-532.

13-Butaye, P., Devriese, L. A. and Haesebrouck, F. (2001). Differences in antibiotic resistance patterns of *Enterococcus faecalis* and *Enterococcus faecium* strains isolated from farm and pet animals. *Antimicrobial Agents and Chemotherapy*, 45: 1374-1378.

14- Cantoni, C., Fabbris, F., and Rogledi, S. (1991). Contenuto in pesticidi organoclorurati in alimenti rilevati nel 1990. *Industria Alimentare*, 30: 383-389.

15-Cantoni, C, and Comi, G. (1997). Changes in the concentrations of pesticide residues in foods and in human tissues between 1960 and 1996. *Outlook on Agriculture*, 26: 47-52.

16-Chaudhury, A. R. Gautam, A. K., and Bhatnagar, V. K. (1990). Lindane induced changes in morphology and lipids profile of testes in rats. *Journal of Biomedical Research*, 49: 1059-1065.

17- Cooper, J., Delahaut, P., Fodey, T. L., and Elliott, C. T. (2004). Development of a rapid screening test for veterinary sedatives and the beta-blocker carazolol in porcine kidney by ELISA. *Analyst*, 129: 169-174.

18- De Brabander, H. F., Batjoens, P., and van Hoof, V. (1992). Determination of thyrostatic drugs by HPTLC with confirmation by GC-MS. *Journal of Planar Chromatography*, 5: 124-130.

19- De Bukanski, B. W., Degroot, J. M., and Beernaert, H. (1988). A two-dimensional highperformance thin-layer chromatographic screening method for sulphonamides in animal tissues. *Zeitschrift für Lebensmittel Untersuchung und Forschung*, 187: 242-245.

20- De Wasch, K., Okerman, L., Croubles, S., De Brabander, H., Van Hoof, J. and De Backer, P. (2001). Detection of residues of tetracycline antibiotics in pork and chicken meat: Correlation between results of screening and confirmatory tests. *Analyst*, 123: 2737-2741.

- 21- Degroodt, J. M., Bukanski, B. W., Beernaert, H. and Courtheyn, D. (1989). Clenbuterol residue analysis by HPLC-HPTLC in urine and animal tissues. *Zeitschrift fur Lebensmittel Untersuchung und Forschung*, 189: 128-131.
- 22- Degroodt, J. M., Bukanski, B. W., De Grood, J., and Beernaert, H. (1991). Cimaterol and clenbuterol residue analysis by HPLC-HPTLC. *Zeitschrift fur Lebensmittel Untersuchung und Forschung*, 192: 430-432.
- 23- Doyle, E. (2000). Human safety of hormone implants used to promote growth in cattle. Food Research Institute, University of Wisconsin, Madison, WI 53706. FRI BRIFFINGS.
- 24- EFSA (2007). Opinion of the scientific panel on contaminants in the food chain on a request from the European Commission related to hormone residues in bovine meat and meat products. *The EFSA Journal*, 510: 1-62.
- 25- Fara, G. M., Del Corvo, G., Bernuzzi, S., Bigatello, A., DiPietro, C., Scaglioni, S., and Chiumello, G. (1979). Epidemic of breast enlargement in an Italian school. *Lancet*, 11: 295-297.
- 26- Franek, M., and Hruska, K. (2005). Antibody based methods for environmental and food analysis: A review. *Veterinary Medicine Czech*, 50: 1-10.
- 27- Gallo, P., Casellano, V., and Serpe, L. (1996). Monitoraggis dei residui di pesticide organochloruratiedorganofosforatiialimenti di origine animale: resultati e consderazioni relative aglianni 1990-1994. *Alimentria* 35: 253-257.
- 28- Garber, E. A. E. (2008). Detection of melamine using comercial enzyme-Linked Immunosorbent Assay technology. *Journal of Food Protection*, 17: 590-594.
- 29- Frenich, G. A., Martinez Vidal, J. L., Cruz Sicilia, A. D., GonzalezRodriguez, M. J. and Plaza Bolanos, P. (2006). Multiresidue analysis of organochlorine and organophosphorus pesticides in muscle of chicken, pork and lamb by gas chromatography-triple quadruple mass spectrometry. *Analytica Chimica Acta*, 558: 42-52.

- 30- Gaudin, V., Cadieus, N., and Maris, P. (2003). Inter-laboratory studies for evaluation of ELISA kits for the detection of chloramphenicol residues in milk and muscles. *Food and Agricultural Immunology*, 15: 143-157.
- 31- Gaugain, M. and Abjean, J. P. (1996). High-performance thin-layer chromaographic method for the fluorescene detection of three nitroimidazole residues in pork and poultry tissue. *Journal of Chromatography A*, 737: 343-346. Gulf Research Center, (2006). Green Gulf Report.
- 32- Gulf Research Center, Dubai. Grundig, B., and Renneberg, R. (2002). Chemical and biochemical sensors. In A. Katerkamp, B. Grundig, and R. Renneberg (Eds.), *Ullmann's encyclopedia of industrial chemistry* (pp. 87-98). Verlag: Wiley-VCH.
- 33- Haagsma, N. (1985). Rapid thin-layer chromatographic screening method for the detection of five sulfonamides in swine tissues: Collaborative study. *Zeitschrift fur Lebensmittel Untersuchung und Forschung*, 181: 45-46.
- 34- Harbottle, H., Thakur, S., Zhao, S., and While, D. G. (2006). Genetics of antimicrobial resistance. *Animal Biotechnology*, 17: 111-124.
- 35- Haughey, S.A., and Baxter, C.A. (2006). Biosensor screening for veterinary drug residues in foddstuffs. *Journal of AOAC International*, 89: 862-897.
- 36- Hernandez, L. M., Fernandez, M. A., Jimenez, B., Gonzalez, M. J., and Garcia, J. F. (1994). Organochlorine pollutants in meats and cow's milk from Madrid (Spain). *Bulletin of Environmental Contamination and Toxicology* 52: 246-253.
- 37- Herrera, A., Arino, A., Conchello, M. P., Lazaro, R., Bayarri, S., and Perez, C. (1994). Organochlorine pesticide residues in Spanish meat products and meat of different species. *Journal of Food Protection*, 57: 441-444.
- 38- Huet, A.C., Mortier, L., Daeseleire, E., Fodey, T., Elliott, C. and Delahaut, P. (2005). Development of an ELISA screening test for nitroimidazoles in egg and chicken muscle. *Analytica Chimica Acta*, 534: 157-162.

- 39- Kadim, I. T., Mahgoub, O., Al-Marzooqi, W., Al-Magbaly, R., Annamal, K. and Khalaf, S. (2010). Enzyme linked immunosorbent assay for screening antibiotic and hormone residues in broiler chicken meat in Sultanate of Oman. *Journal of Muscle Foods*. 21(2): 243 – 254.
- 40- Lee, H. J., Lee, M. H., Ryu, P. D., Lee, H. and Cho, M. H. (2001). Enzyme-linked immunosorbent assay for screening the plasma residues of tetracycline antibiotics in pigs. *Journal of Veterinary Medicine*, 63: 553-556.
- 41- Lembowicz, K., Sitarska, E., Gorski, T., and Ludwicki, J. K. (1991). The effect of organic chlorine compounds and their metabolites present in human milk on newborn mice. *Toxicology Letters*, 57: 215-226.
- 42- Lone, K. P. (1997). Natural sex steroids and their xenobiotics analogs in animal production: growth, carcass quality, pharmacokinetics, metabolism, mode of action, residues, methods and epidemiology. *Critical Reviews in Food Science and Nutrition*, 37: 93-209.
- 43- MacLachlan, D. J. and Bhula, R. (2008). Estimating the residue transfer of pesticides in animal feedstuffs to livestock tissues, milk and eggs: a review. *Australian Journal of Experimental Agriculture*, 48: 589-598.
- 44- Magana, J. A. and Calderon de la Barca, A. M. (2008). Risk assessment of genetically modified crops for nutrition and health. *Nutrition Reviews*, 67: 1-16.
- 45- Mahgoub, O., Kadim, I. T., Ann Mothershaw, Al Zadjali, S. A., and Annamalai, K. (2006). Use of Enzyme Linked ImmunoSorbent Assay (ELISA) for detection of antibiotic and anabolic residues in goat and sheep meat. *World Journal of Agricultural Sciences*, 2: 298-302.
- 46- McEvoy, J. D. (2002). Contamination of animal feedstuffs as a cause of residues in food: A review of regulatory aspects, induce and control. *Analytica Chimica Acta*, 473: 3-26.
- 47- McEwen, S. A. (2006). Antibiotic use in animal agriculture: what have we learned and where are we going?. *Animal Biotechnology*, 17: 239-250.

McEwen, S. A. and Fedorka-Cray, P. (2002). Antimicrobial use and resistance in animals. *Clinical Infection Disease*, 34: 93-106.