

**Republic of Iraq
Ministry of Higher Education
& Scientific Research
University of Al-Qadisiyah
College of Veterinary Medicine**



**Jaagsiekte Sheep Retrovirus (JSRV)
from virus to lung cancer in sheep**

A Graduation Project Submitted to the Department
Council of the Internal and Preventive Medicine-College
of Veterinary Medicine/ University of Al-Qadisiyah in a
partial fulfillment of the requirements for the Degree of
Bachelor of Science in Veterinary Medicine and Surgery.

By
Ali shamki Idan and Hussain Kazem Jawed

Supervised by
Asst.Prof. Hala Abbas Naji

2021 A.D.

1441 A.H.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

فَتَعَلَى اللَّهِ الْمَلِكُ الْحَقُّ وَلَا تَعْجَلْ بِالْقُرْآنِ مِنْ قَبْلِ أَنْ يُقْضَىٰ
إِلَيْكَ وَحْيُهُ، وَقُلْ رَبِّ زِدْنِي عِلْمًا ﴿١١٤﴾

صَدَقَ اللَّهُ الْعَظِيمُ،

من سورة طه

Certificate of Supervisor

I certify that the project entitled (Jaagsiekte Sheep Retrovirus (JSRV) from virus to lung cancer in sheep) was prepared by **Ali shamki Idan and Hussain Kazem Jawed** under my supervision at the College of Veterinary Medicine / University of Al-Qadisiyah.

Supervisor

Asst.Prof. Hala Abbas Naji

Dept. of Pathology &Poultry disease

Coll. Of Vet.Med./ Univ. of Al-Qadisiyah.

-- / -- / 2021

Certificate of Department

We certify that Ali shamki Idan and Hussain Kazem Jawed had finished their Graduation Project entitled (Jaagsiekte Sheep Retrovirus (JSRV) from virus to lung cancer in sheep) and candidate it for debating.

Instructor

Dr. Muthanna H. Hussain

-- / -- / 2021

Head of Dept of Int. and Prev. Med.

Dr.asst.prof. Saed H. Al-Hussiny

-- / -- / 2021

Abstract

Jaagsiekte Sheep Retrovirus (JSRV) is a betaretrovirus infecting sheep. This virus is re-sponsible for a pulmonary adenocarcinoma, by transformation of epithelial cells from the bronchioli and alveoli. This animal cancer is similar to human bronchioloalveolar cancer (BAC), a specific form of human lung cancer for which a viral aetiology has not yet been identified. JSRV interacts with target cells through the membrane receptor Hyal2. The JSRV genome is simple and contains no recognised oncogene. It is now well established that the viral envelope protein is oncogenic by itself, via the cytoplasmic domain of the transmembrane glycoprotein and some domains of the surface glycoprotein. Activation of the PI3K/Akt and MAPK pathways participates in the envelope-induced transformation. Tumour development is associated with telomerase activation. This review will focus on the induction of cancer by JSRV.

Table of contents

INTRODUCTION	3
OPA: A VIRUS-INDUCED CANCER	4
Natural history	
Pathogenesis	4
JSRV: a member of the Betaretrovirus genus	5
Virus and genomic organisation	5
Virus and genomic organisation	9
CONCLUSION	13
REFERENCES	14

INTRODUCTION

Ovine pulmonary adenocarcinoma (OPA) is a contagious tumour originating in the distal lung after infection by Jaagsiekte Sheep Retrovirus (JSRV). The disease was initially described in 1915 in South Africa and is present worldwide. JSRV belongs to the *Retroviridae* family and the *Betaretrovirus* genus. OPA is similar to a peculiar form of human cancer, bronchioloalveolar cancer (BAC), with which it shares clinical, radiological and histopathological features [51]. Although the molecular mechanisms of JSRV induced-tumorigenesis are still only partially understood, the significant recent efforts in the field of OPA research reinforce the for.

JSRV is the virus that is the cause of the contagious lung tumors in sheep called ovine pulmonary adenocarcinoma (OPA). The disease has also been called "jaagsiekte", after the Afrikaans words for "chase" (jaag) and "sickness" (siekte), to describe the respiratory distress observed in an animal out of breath from being chased, indicating the breathing difficulty experienced by infected sheep. Transmission of virus is through aerosol spread between sheep.

The exogenous infectious form of JSRV has an endogenous counterpart which is present in the genomes of all sheep and goats.[1] The sheep genome has around 27 copies of endogenous retroviruses (enJSRVs) that are closely related to JSRV. Endogenous JSRV has several roles in the evolution of the domestic sheep as they are able to block the JSRV replication cycle and play a critical role in sheep conceptus development and placental morphogenesis.[2]

Although OPA resembles human lung cancer, human lung cancer is not known to be caused by betaretroviruses.[3] Even though a possibility of a viral cause has been eliminated in bronchoalveolar cancer, understanding the molecular mechanisms leading to the transformation of lung epithelia by JSRV may be of interest in the context of therapeutic approaches in human lung cancers in general and bronchoalveolar adenocarcinoma (BAC) in particular.[

OPA: A VIRUS-INDUCED CANCER

Natural history

Transmission of OPA among sheep has been suspected for almost two centuries. The first report dates back to 1825, with a letter written by a farmer who complained about the loss of many of his sheep. The disease was called “Jaagsiekte”, after the Afrikaans words for “chase” (Jaag) and “sickness” (siekte), to describe the respiratory distress observed in an animal out of breath from being chased [2]. The first evidence of a viral cause came from the observation of retrovirus particles in the lungs from sheep presenting clinical signs of cancer [1], and was further clearly confirmed by experimental induction of the disease by intratracheal inoculation of viral particles with a reverse transcriptase activity [4], cytoplasmic fractions of tumoral cells [2-1] or pulmonary secretions [1].

The disease can also be efficiently transmitted to goats by experimental inoculation [3-1]. JSRV has been definitively demonstrated as the aetiological agent of OPA by experimental inoculation of particles produced from a JSRV-molecular clone [1].

OPA is present on all continents. Its incidence is difficult to evaluate in the absence of an appropriate screening tool. The virus is transmitted between animals by close contact, mainly through aerosolized particles. The breeding conditions are of major importance for the dissemination of the virus. The incubation period in naturally infected animals ranges between 2 and 4 years [3], but the cancer may be diagnosed as early as a few months after birth. The incubation period may vary according to the type of infection (experimental versus spontaneous infection), and the age of the animals [5]. Interestingly, injection of tumoural tissues into newborn lambs rapidly induces the disease in 3–6 weeks [7]. In natural conditions, the rapid development of tumoral lesions in young animals suggests a greater susceptibility of the developing lung to JSRV [8]. In utero transmission to the fetus has been suggested [8].

Pathogenesis

JSRV is transmitted by the respiratory route and may also infect lymphocytes and myeloid cells, in addition to the lung epithelia. Expression of the JSRV Envelope protein activates signalling cascades that promote cellular proliferation and malignant transformation of the cells. Initially, the tumour cells grow along the alveolar walls in a pattern reminiscent of human BAC, but subsequently become more invasive and metastasize to the local lymph nodes. Larger tumours may be necrotic and fibromatous at their centre. As the tumour grows, fluid production in the lung increases and this is likely to promote virus spread to other sheep. Only when the tumour reaches a size large enough to compromise lung function, do clinical signs appear. Critically, the majority of infected animals in endemic areas never show outward signs of infection, but they may be shedding virus, thus promoting inadvertent introduction of the disease into previously unaffected flocks and new geographical areas

JSRV: a member of the *Betaretrovirus* genus

JSRV is the virus responsible for the induction of OPA. JSRV belongs to the family *Retroviridae*, to the subfamily *Orthoretrovirinae* and the genus *Betaretrovirus*. The *Betaretrovirus* genus also comprises Mouse Mammary Tumour Virus (MMTV) responsible for a mammary adenocarcinoma in the mouse, Mason-Pfizer Monkey Virus (MPMV) isolated from a Rhesus monkey, and Squirrel Monkey RetroVirus (SMRV)(7). In 2003, Dolly the sheep, the first mammal cloned from an adult cell, died after being diagnosed with JSRV and lung cancer in sheep(3)

Virus and genomic organisation

Retroviruses are RNA viruses infecting vertebrate species and many non-vertebrates. Virions (80–100 nm in diameter) are spherical and surrounded by an envelope, with spikes composed of virus-encoded glycoproteins. The envelope is composed of viral proteins and elements of the host-cell membrane (lipid bilayer and proteins). The virions carry two copies of the genome, composed of linear, positive, single-stranded RNA. The 7.58 kb genome of the infectious virus comprises four main genes, organised as 5'- *gag*- *pro*- *pol*- *env* - 3' (Fig. 1), which encode the virus proteins (reviewed in [4-6]). Interestingly, while the nucleotide sequences of *gag*, *pro* and *pol* are homologous to their counterparts in MPMV, the *env* gene of JSRV is more related to that of MMTV and Human Endogenous Retrovirus -K (HERV-K). JSRV is organised as a simple retrovirus, with an additional open reading frame, named ORF-

x, that overlaps the 3' end of the *pol* gene. ORF-x is unique among retroviruses and may encode a putative accessory protein of 166 amino acids. Although the existence of the protein is debated, ORF-x is strongly conserved in various isolates, and sequence analyses suggest a selective pressure for its conservation [6- 8] (and our unpublished data). Interestingly, two sub-genomic mRNA with splice acceptor sites within or in the vicinity of ORF-x have been identified, suggesting that this putative gene may actually be transcribed [3]. We have also identified these two mRNA in tumoral lungs (unpublished data).

The JRSV genome contains non-coding regions at the ends of the genome, that are essential for the virus replication: R repeated at both ends; U5 unique to the 5' end and U3 unique to the 3' end. The integrated genome is flanked by the long terminal repeat (LTR) composed of the 3 regions U3-R-U5(2). The LTR serve as the sites of transcriptional initiation. The U3 region contains several elements important for viral transcription and tropism. The *gag* gene (*group-specific- antigen*) encodes a single polyprotein that is cleaved into at least three proteins: the matrix (MA), the major capsid protein (CA) and the nucleocapsid (NC). The *pro* gene encodes a protein compatible with already-described dUTPase in its 5' part and a protease (PRO) in its 3' end. Whereas the protease cleaves the precursor polyproteins, the dUTPase (reviewed in [8]) prevents the incorporation of deoxyuridine triphosphate (dUTP) by the reverse transcriptase [1]. The *pol* gene is predicted to encode the enzymatic activities: Reverse Transcriptase (RT) and integrase (IN), respectively implicated in the replication of the viral RNA and in the integration of the retrotranscribed DNA provirus into the host genome. The RT is a RNA-dependent DNA polymerase, essential for the conversion of viral RNA into DNA. The *env* gene encodes the surface (SU) and transmembrane (TM) glycoproteins. The SU glycoprotein interacts with the cellular receptor of JSRV in sheep, Hyal-2, a glycosylphosphatidylinositol anchored hyaluronidase-2 [7]. The TM glycoproteins presumably anchor SU into the

lipid bilayer and are composed of a hydrophobic stretch of amino acids followed by a short cytoplasmic tail. The JSRV envelope is a major determinant for the cellular transformation as discussed in detail below.

JSRV is phylogenetically related to the Enzootic Nasal Tumour Virus (ENTV), the agent responsible for nasal adenocarcinoma, a contagious tumour of the mucosal nasal glands affecting sheep and goats. In infected animals, epithelial cell proliferation is responsible for continuous nasal discharge, respiratory distress, exophthalmos and important skull deformations (for review see [8]). Co-infection of ENTV and JSRV has been reported [4,13].

A family of endogenous retroviruses, enJSRV (endogenous JSRV) closely related to JSRV, is present in domestic and wild sheep and goats [3-4]. JSRV and en-JSRV genomes are highly related with 90–98% homology in deduced amino-acid sequences [5]. Endogenous retroviruses (ERV) are vertically transmitted as stable Mendelian genes in the germline of most eukaryotes. They derive from the integration of exogenous viruses in the host genomes, followed by genetic stabilisation through accumulation of mutations. Several families of HERV such as HERV-F, HERV-FRD, HERV-K, HERV-R, HERV-T and HERV-W (defined by the tRNA complementary to their putative primer binding site using the one letter code for the tRNA's corresponding amino acid) have been described in the human genome. Eight to twelve copies of enJSRV have been located on metaphase chromosome spreads of sheep and goats using fluorescent in situ hybridisation [8,14] (Fig. 2).

The biological significance of ERV is still largely debated. Expression of ERV has been described in the placenta and genital tract of mammals including humans with HERV-W. In the female genital tract, enJSRV expression is limited to the epithelia and is particularly abundant in the




Figure 1. Subtypes of lesions observed in OPA: bronchioloalveolar (A), acinar (B) and papillary (C). (A color version of this figure is available at www.edpsciences.org/vetres.)

wheelbarrow test”) is the flow of abundant mucoid fluid (up to 500 mL per day in advanced disease) from nostrils when the rear legs of the animal are raised. Extrathoracic metastases are rare. Macroscopic examination shows enlarged lungs infiltrated with tumoural areas, varying from small nodules (1–30 mm) to lobar consolidation [8,12]. The disease is multifocal, disseminated in both lungs, pneumonic in appearance, with the airways filled with fluid produced by the tumoural cells.

According to the latest WHO classification for human lung cancers [7], OPA should be referred to as a mixed adenocarcinoma with associated bronchioloalveolar, papillary and/or acinar subtypes. The bronchioloalveolar differentiation is characterised by the expansion of cells following the alveolar septa, also referred to as the lepidic spread, without destruction of the alveolar architecture. The papillary subtype is defined by the presence of papilla-like structures protruding above the epithelial layer and replacing the underlying alveolar architecture. Finally, the acinar subtype is composed of duct-like structures or acini and tubules composed of cells resembling bronchial glands. In JSRV-induced adenocarcinoma, these three histopathological subtypes may be present within the same tumoural tissue(6,10).

The OPA tumoural cells derive from epithelial cells of the distal lung, namely alveolar type II cells (for ~80% of the cells) and Clara cells (for ~15% of the cells) [7]. Type II pneumocytes produce different components of surfactant, a tensio-active agent that allows the maintenance of alveolar integrity.

Interestingly, tumoural cells over-express CD208/DC-LAMP (Dendritic Cell Lysosomal Associated Membrane Protein) [4], a protein belonging to the LAMP family of proteins (Lysosomal-Associated Membrane Protein), and initially described in activated dendritic cells. CD208/DC-LAMP is also constitutively expressed in human, murine and ovine type II pneumocytes, and is over-expressed in human bronchioloalveolar carcinoma and OPA [7]. In type II pneumocytes, CD208 is expressed in the constitutive membranes of the lamellar bodies, specific vesicles specialised in surfactant production. The role of CD208 over-expression in tumoural development is still unknown.

Mechanisms of oncogenesis

Oncogenic retroviruses have been historically divided into two groups




Figure 2. Microscopic pattern of lesions from OPA. Typical aspect of a non-tumoural (A) and tumoural ovine lung (B) labelled with an anti SP-A (Surfactant Protein A) antibody. Arrows show tumoural cells, i.e. type II pneumocytes labelled with an anti SP-A antibody. (A color version of this figure is available at www.edpsciences.org/vetres.)

depending on the speed of disease induction. Acute and non acute retroviruses cause tumours with a short or long latency period respectively. Acute retroviruses have been described in mice, birds, primates and are oncogene-bearing retroviruses. Viral oncogenes are derived from normal cellular genes or proto-oncogenes, involved in regulation of cell proliferation. Once

captured by the virus, proto-oncogenes undergo mutations that lead to uncontrolled cell proliferation. Rous Sarcoma Virus (RSV) is an example of avian retrovirus bearing an oncogene, *v-src*, deriving from a mammalian cellular kinase.

Typical lesions of OPA may be observed in just a few weeks following experimental inoculation of concentrated JSRV particles [6,4, 8]. This timeframe of disease induction and the multifocal pattern of OPA are compatible with the presence of an oncogene in the JSRV genome [8]. JSRV DNA transfection in murine NIH 3T3 fibroblasts induces foci of transformed cells, clearly indicating that the JSRV genome contains an oncogenic element [4, 7]. However, to date no sequence homologous to

any cellular oncogene has been detected in the JSRV genome [7,9]. The open reading frame (orf-x) of JSRV might be a potential oncogene, although it does

not present any sequence homology with known oncogenes [6, 7], and its deletion does not prevent transformation in vitro.

It is now clearly established that JSRV induces tumours via the oncogenic properties of its envelope, which is both necessary and sufficient to induce transformation. The transforming property of the JSRV envelope has been demonstrated in vitro in various cell lines including murine NIH 3T3 fibroblasts [4], rat 208F fibroblasts [7], avian DF-1 fibroblasts [4, 3], bronchial human BEAS-2B epithelial cells [1], canine kidney MDCK epithelial cells [3], and rat kidney RK3E cells [6]. The oncogenic property of the JSRV envelope has also been shown in vivo in an immunodeficient mouse model [8] and recently in sheep [7]. Expression of the JSRV envelope in mice using a replication-incompetent adeno-associated virus vector, AAV6, results in lung tumours similar to those observed in sheep [7]. The bronchioloalveolar localisation of the tumours and the expression of the surfactant protein SP-C show that the transformed cells are derived from type II pneumocytes [8]. Using a replication-defective virus carrying the *env* gene under the control of the JSRV LTR, it has been shown that the JSRV envelope is sufficient to induce lung tumours in sheep. The envelope of the related ENTV virus displays the same oncogenic property both in vitro [2, 3] and in vivo [8,11].

Deletion experiments show that the TM (transmembrane) region of the envelope is the main determinant for cell transformation [1, 3, 8]. In addition to the TM region, deletions of the SU (surface) glycoprotein (going from the signal peptide to the junction between the SU and TM subunits) also abolish transformation induced by the envelope, suggesting that multiple domains of SU may be involved in cellular transformation. The cytoplasmic tail of TM, composed of 43 amino acids, is essential for the transformation process in MDCK and NIH-3T3 cells [4, 6,12]. Corresponding to a potential consensus site (phosphorylated on tyrosine Y) linked to the SH2 domain of the p85 subunit of PI3K (Phosphatidylinositol-3 Kinase), a kinase that activates Akt. The PI3K-Akt signalling pathway is determinant in cellular proliferation and survival (for review [1]). Following a membrane stimulus such as a growth factor binding to its receptor, PI3K is recruited to the cell membrane. Phosphorylated-PI3K phosphorylates the second messenger PIP2 (phosphatidylinositol (4,5) biphosphate) into active PIP3 (phosphatidylinositol (3,4,5,13) triphosphate). PIP3 can then recruit PDK1 (phosphatidylinositol-dependent kinase 1), which in turn phosphorylates Akt. The kinase Akt can phosphorylate diverse substrates involved in signalling cascades controlling cellular proliferation, survival and metabolism. Akt phosphorylation inhibits proteins such as GSK-3 (Glycogen Synthase Kinase 3), FOXO (forkhead box

transcription factor), p24 or Kip1, and activates mTOR (mammalian target of rapamycin), an important regulator of cell growth.

The mechanisms leading to JSRV-induced cell transformation are in fact much more complex than previously considered. Several experiments rule out a direct role for the YXXM motif in Akt activation. In JSRV-transformed cells, phosphorylation of the Y590 residue of the intra-cytoplasmic TM tail, or interaction between the p85 subunit of PI3K and the envelope (a prerequisite for Akt activation), have never been shown [4, 3]. Mutations of the YXXM motif do not abolish Akt phosphorylation, so that the exact role of this motif remains to be determined [2, 4, 6]. Moreover, implication of the YXXM motif may differ between cell types in in vitro experiments [4, 6, 1]; while it is essential for transformation of NIH 3T3 fibroblasts, it is not required for transformation of avian DF-1 fibroblasts. Nevertheless, the presence of the YXXM motif seems to affect the efficiency of envelope-mediated transformation [3]. Since the YXXM motif does not explain the oncogenic properties of JSRV, the mechanism of Akt activation in transformed cells still need to be determined. Interestingly, treatment of cells with LY294002, a PI3K-specific inhibitor, drastically reduces Akt phosphorylation in NIH 3T3 cells [2,5,11,14], suggesting that PI3K-dependent Akt activation may occur.

CONCLUSION

Over the last few years, research on JSRV and the resulting OPA has focussed on the molecular mechanisms of cell trans-formation. Interesting results came from the discovery that the virus envelope acts as a potent oncogene. Several cellular pathways seem to occur in the cell trans-formation.

To conclude, research into the biology of JSRV and mechanisms leading to the development of OPA is of great interest both for the naturally induced cancer in sheep and for BAC, the related human cancer. Even though a viral agent remains uncertain in BAC patients, understanding the steps leading to the transformation of lung epithelia may be of interest in the context of therapeutic approaches in human lung cancers in general and BAC in particular.

REFERENCES

- [1] Aizawa S., Suda Y., Furuta Y., Yagi T., Takeda N., Watanabe N., Nagayoshi M., Ikawa Y., Env-derived gp55 gene of Friend spleen focus-forming virus specifically induces neoplastic proliferation of erythroid progenitor cells, *EMBO J.* (1990) 9:2107–2116.
- [2] Alberti A., Murgia C., Liu S.L., Mura M., Cousens C., Sharp M., Miller A.D., Palmarini M., Envelope-induced cell transformation by ovine betaretroviruses, *J. Virol.* (2002) 76:5387–5394.
- [3] Alian A., Sela-Donenfeld D., Panet A., Eldor A., Avian hemangioma retrovirus induces cell proliferation via the envelope (*env*) gene, *Virology* (2000) 276:161–168.
- [4] Allen T.E., Sherrill K.J., Crispell S.M., Perrott M.R., Carlson J.O., DeMartini J.C., The jaagsiekte sheep retrovirus envelope gene induces transformation of the avian fibroblast cell line DF-1 but does not require a conserved SH2 binding domain, *J. Gen. Virol.* (2002) 83:2733–2742.
- [5] Archer F., Jacquier E., Lyon M., Chastang J., Cottin V., Mornex J.-F., Leroux C., Alveolar Type II cells isolated from pulmonary adenocarcinoma: a model for JSRV expression in vitro, *Am. J. Respir. Cell. Mol. Biol.* (2006) (in press).
- [6] Bai J., Bishop J.V., Carlson J.O., DeMartini J.C., Sequence comparison of JSRV with endogenous proviruses: envelope genotypes and a novel ORF with similarity to a G-protein-coupled receptor, *Virology* (1999) 258:333–343.
- [7] Beasley M.B., Brambilla E., Travis W.D., The 2004 World Health Organization classification of lung tumors, *Semin. Roentgenol.* (2005) 40:90–97.
- [8] Caporale M., Centorame P., Giovannini A., Sacchini F., Di Ventura M., De las Heras M., Palmarini M., Infection of lung epithelial cells and induction of pulmonary adenocarcinoma is not the most common outcome of naturally occurring JSRV infection during the commercial lifespan of sheep, *Virology* (2005)
- [9]

York DF, Querat G (2003). "A history of ovine pulmonary adenocarcinoma (jaagsiekte) and experiments leading to the deduction of the JSRV nucleotide sequence". *Current Topics in Microbiology and Immunology*. 275: 1–23.

[10]

Leroux C, Girard N, Cottin V, Greenland T, Mornex JF, Archer F (2007). "Jaagsiekte Sheep Retrovirus (JSRV): from virus to lung cancer in sheep". *Veterinary Research*. 38 (2): 211–28. doi:10.1051/vetres:2006060. PMID 17257570.

[11]

Griffiths DJ, Martineau HM, Cousens C (May 2010). "Pathology and pathogenesis of ovine pulmonary adenocarcinoma". *Journal of Comparative Pathology*. 142 (4): 260–83. doi:10.1016/j.jcpa.2009.12.013. PMID 20163805

[12]

Hopwood P, Wallace WA, Cousens C, Dewar P, Muldoon M, Norval M, Griffiths DJ (November 2010). "Absence of markers of betaretrovirus infection in human pulmonary adenocarcinoma". *Human Pathology*. 41 (11): 1631–40.

[13]

Varela M, Spencer TE, Palmarini M, Arnaud F (October 2009). "Friendly viruses: the special relationship between endogenous retroviruses and their host". *Annals of the New York Academy of Sciences*. 1178: 157–72. doi:10.1111/j.1749-6632.2009.05002.x. PMC 4199234. PMID 19845636.

[14]

Armezzani A, Varela M, Spencer TE, Palmarini M, Arnaud F (December 2014). "'Ménage à Trois': the evolutionary interplay between JSRV, enJSRVs and domestic sheep". *Viruses*. 6 (12): 4926–45. doi:10.3390/v6124926. PMC 4276937. PMID 25502326.

الخلاصة

هو فيروس بيتاريتروفيروس يصيب الأغنام. يمكن إعادة (JSRV) Jaagsiekte Sheep Retrovirus - رعاية هذا الفيروس لسرطان غدي رئوي ، عن طريق تحويل الخلايا الظهارية من القصيبات والحوبيصلات ، وهو شكل محدد من سرطان (BAC) الهوائية. يشبه سرطان الحيوان هذا سرطان القصيبات السنخي البشري مع الخلايا المستهدفة من خلال مستقبل الغشاء JSRV الرئة البشري لم يتم تحديد مسبباته الفيروسية بعد. يتفاعل بسيط ولا يحتوي على أي مادة أورام معترف بها. لقد ثبت الآن جيداً أن بروتين JSRV إن جينوم Hyal2. الغلاف الفيروسي هو منشئ الورم في حد ذاته ، عبر المجال السيتوبلازمي للبروتين السكري عبر الغشاء يشارك في التحول MAPK و PI3K / Akt وبعض مجالات البروتين السكري السطحي. تنشيط مسارات الناجم عن المغلف. يرتبط تطور الورم بتنشيط التيلوميراز. ستركز هذه المراجعة على تحريض السرطان بواسطة JSRV